VIEQUES

Dr. Arturo Massol Deyá



Departamento de Biología Universidad de Puerto Rico Mayagüez, PR 00681-9012



1940-2003

Elements of a Bomb

Acetone

- * Acetyl Triethyl Citrate
- * Aluminum Powder
- * Ammonium Nitrate

Antimony

- **Barium**
- * Boron
- * Butyl Stearate

Cadmium

Chromium

- * Diazodinitrophenol Diethyl Phthalate Dimethyl Phthalate Di-n-butyl phthalate Di-n-octyl Phthalate
- * Dichromated Aluminum Powder

- 2, 4- Dinitrotoluene
- 2, 6-Dinitrotoluene
- Diphenylamine
- * Ethyl Centralite
- * HMX

Lead

* Magnesium Powder Magnesium Alloy Powder


Mercury

- **Nickel Powder**
- Nitrocellulose
- * Nitroglycerin
- * Nitroguanidine
- * Nitrostarch
- * Pentolite
- * Pentaerythritol Tetranitrate (PETN)
- * Phosphorous


- * Potassium Nitrate
- * RDX (Cyclonite)
- * Resorcinol Selenium
- * Sodium Nitrate
- * Strontium Nitrate
- * Tetracene
- * Tetryl
- * Titanium Powder
- * TNR (Trinitroresorcinol)
- * TNT(Trinitotoluene)
- * Tri Amino Guanidine Nitrate
 - Vinyl Acetate
 - Vinyl Chloride
 - Zinc
- * Zirconium Powder









Dispersion of natural trace elements is increased due to military activities. Is pollution at the Atlantic Fleet Weapons Training Facility contained?









Why PLANTS?

- Transitory receptors of trace elements.
- Plants can concentrate environmental toxins, thus they can be good bioindicators of environmental degradation.


 Ecological significance: start-up of the food chain.
 Plants are relatively "immune" to human toxins. (no diabetes, hypertension or cancer cases have been reported in plants)

Potential use in clean up efforts: Phytoremediation



Methodology


- Life vegetation samples were manually collected.
- Each station was randomly sampled in a 2 to 5 m² circular plot. Sampling sites were independently established for each species. Vegetation analysis always included leaves material. Other plant components were selected as indicated. The sampling was consistently performed by selecting only large, green, healthy looking leaves. Samples from each species consisted of over 30 leaves picked alternately from upper, middle, and the lower sections of 5 to 10 individual plants.
- Samples were acid digested (2-4 replicates) and analyzed in an atomic absorption spectrophotometer.




Autoridad de Carreteras y Transportación Oficina de Fotogrametría Fecha: 27 de febrero de 1996 Escala Aproximada: 1:20,000 Area: Punta Este, Vieques (Puerto Rico)

Areas de Muestreo:

- Monte David
- 🛛 Playa Carrucho
- 🛛 Noroeste de la Laguna Gato
- Laguna Icacos







Zonas Bajo Estudio por Massol y Díaz. CasaPueblo, Adjuntas PR.


Areas de Muestreo


- (a) Polígono de Prácticas Militares, Vieques
- (b) Zona Civil, Vieques
- **c)** Antigua Zona de Prácticas Militares, Culebra
- **d)** Bosque Xerofítico de Guánica
- e Fincas, Mayagüez y Las Marías
- **f**) Zona Industrial, Peñuelas











...analysis revealed sea around Vieques is free of heavy metal contamination!

However, at that time, live However, at that time, live ammunition wasn't being used.

Government Documentation

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY REGION II

IN THE MATTER OF:

United States Department of the Navy

ADMINISTRATIVE ORDER ON CONSENT

DOCKET No. RCRA-02-2000-7301 Proceeding under Section 3008(h),

of the Resource Conservation and

Recovery Act, as amended.

Atlantic Fleet Weapons Training Facility Viegues Island, PR

EPA ID Number PRD980536221

Respondent


In 1994, the EPA identified 102
 violations to water quality parameters.

 The EPA Discharge Monitoring Reports identified excessive concentrations of the following contaminants in the coastal waters of Vieques:

Boron Grease Phenols Selenium Silver Sulfates Zinc Cyanide (10x) Cadmium (240x) Chromium (13x) Iron Lead (105x) Manganese Mercury (4.6x) Arsenic (6.6x) Nitrogen (188x)

	Chemical Element (µg/g DW)							
	Pb	Co	Ni	Al	As	Cd	Cu	
AFWIF-2001 (Active)	33.32 (10.77)	29.60 (5.51)	2866 (1.58)	-	-	0.28 (0.13)	30.48 (3.63)	
AFW'IF-2004(Non-active)	8.14 (3.15)	1061 (4.21)	3.43 (2.13)	1543 (67.7)	0.61 (0.46)	0.15 (0.15)	17.42 (1.83)	
Guțnica-2001	5.58 (1.90)	4.19 (0.06)	1464 (4.75)	-	-	0.28 (0.01)	15.39 (4.16)	
Guțnica-20032004	2.33 (2.24)	1.82 (0.43)	3.75 (1.85)	3412 (112.9)	1.04 (0.25)	0.28 (0.22)	1216 (2.83)	





Pigeon Peas

Leaf samples (June 2000)

	Cu	Mn	Zn	Pb	Ni	Cd	Со
Monte Carmelo/Vieques	18.30	182.99	34.77	10.98	10.37	2.44	14.64
	12.58	203.73	27.56	10.19	10.19	3.00	14.98
average	15.44	193.36	31.17	10.59	10.28	2.72	14.81
std dev	(4.04)	(14.67)	(5.10)	(0.56)	(0.13)	(0.40)	(0.24)
Sector Gobeo/Vieques	28.87	77.63	87.57	54.43	7.10	0.95	7.10
	22.75	71.56	71.56	43.85	6.20	1.24	4.96
average	25.81	74.60	79.57	49.14	6.65	1.10	6.03
std dev	(4.33)	(4.29)	(11.32)	(7.48)	(0.64)	(0.21)	(1.51)
CONTROL	10.45	142.98	31.62	4.95	7.97	0.82	4.40
Cerro Las	11.33	132.69	29.58	4.98	7.46	0.83	3.87
Mesas/Mayagüez	11.08	130.69	30.4	4.55	7.67	0.85	4.55
	11.30	143.36	31.98	5.24	6.66	1.10	4.41
promedio	11.04	137.43	30.90	4.93	7.44	0.90	4.31
std dev	(0.41)	(6.68)	(1.11)	(0.28)	(0.56)	(0.13)	(0.30)
MC VIEQ/CONTROL	1.4	1.4	1.0	2.1	1.4	3.0	3.4
SG VIEQ/CONTROL	2.3	0.5	2.6	10.0	0.9	1.2	1.4



Monte Carmelo, Vieques (Puerto Rico) Pigeon Peas


	Cu	Zn	Ni	Pb	Cd	Со		
Leaf samples	15.00	39.15	2.45	14.70	2.58	16.78		
	6.43	25.08	2.06	15.53	2.26	18.13		
average	10.72	32.12	2.26	15.12	2.42	17.46		
std dev	(6.06)	(9.95)	(0.28)	(0.59)	(0.23)	(0.95)		
Fruit samples	7.02	35.31	5.33	7.80	2.21	23.29		
	9.41	32.68	4.27	4.89	4.13	28.21		
average	8.22	34.00	4.80	6.35	3.17	25.75		
std dev	(1.69)	(1.86)	(0.75)	(2.06)	(1.36)	(3.48)		
FRUIT/LEAF	0.8	1.1	2.1	0.4	1.3	1.5		
					August 2001			

2ND Draft Preliminary Evaluation of Health Risks Related to Naval Activities at the Atlantic Fleet Weapons Training Facility on Vieques, Puerto Rico

August 2000

"Bombing activities that occur on the eastern portion of the island are likely to produce airborne emissions that would be carried in a westerly direction." US Navy

20,000 - 30,000 explosion every year! > 50 years!












QuickTime™ and a TIFF (LZW) decompressor are needed to see this picture.



Caribbean Journal of Science, Vol. 39, No. 2, 000–000, 2003 Copyright 2003 College of Arts and Sciences University of Puerto Rico, Mayagüez

Trace Element Composition in Forage Samples from a Military Target Range, Three Agricultural Areas, and One Natural Area in Puerto Rico


Elba Díaz and Arturo Massol-Deya

Department of Biology, University of Puerto Rico, Mayagüez, Puerto Rico 00681 amassol@stahl.uprm.edu

ABSTRACT scarce. Trace Facilities (Al State Forest. with acid. C Rico. Simila Sporobolus a and marine

QuickTime[™] and a TIFF (LZW) decompressor are needed to see this picture.





Agradecimientos

»Elba Díaz
»Dustin Pérez
»Ernie Pérez
»Manuel Berríos

- Botánicos [UPR-Mayagüez]
- Familia Zenón-Encarnación
- Programa de Biotecnología Industrial, UPRM
- Casa Pueblo, Adjuntas
- Departamento de Salud de PR
- Escuela Elemental Playa Grande, Victor Belardo y los estudiantes de programa ambiental